



# Exploring the Therapeutic Potential of Amino Acids in the Treatment of Psychoactive Substance Addiction: A Comprehensive Review

Kian Zehtabian<sup>1</sup>, Dennis Relajo-Howell<sup>2,3\*</sup>, Ronald C. Yrog-irog<sup>4</sup>, Aidin Kazemizadeh<sup>5</sup>

<sup>1</sup>Islamic Azad University, Iran.

<sup>2</sup>Psychreg, United Kingdom.

<sup>3</sup>University of Edinburgh, United Kingdom.

<sup>4</sup>Cebu Institute of Technology-University, Philippines.

<sup>5</sup>King's College London, United Kingdom.

## Abstract

*This paper offers a comprehensive review of the potential therapeutic role of amino acids, specifically tyrosine and tryptophan, in treating addiction to psychoactive substances such as cocaine and amphetamine. Amino acids are fundamental to various physiological processes and serve as precursors to neurotransmitters like dopamine. This review hypothesises that amino acids could influence dopamine production in the brain, offering a novel treatment pathway for addiction. While preliminary human trials show promise, the exact mechanisms and potential side effects are not yet fully understood, particularly in the context of co-occurring mental health disorders.*

**Keywords:** Addiction, Amino Acids, Dopamine, Mental Health, Psychoactive Substances

## INTRODUCTION

Amino acids serve as essential biomolecules, comprising amino groups ( $-NH_3^+$ ), carboxylate groups ( $-COO^-$ ), and a side chain ( $-R$ ). Of the hundreds of amino acids identified or synthesised, only 20 commonly serve as the building blocks of proteins. With the exception of glycine, all  $\alpha$ -amino acids are chiral molecules, featuring four distinct substituents attached to the  $\alpha$ -carbon (Maloy & Hughes, 2013). Beyond their role as protein subunits, amino acids also function in osmoregulation, neurotransmission, and metabolic processes (Bhagavan & Ha, 2015).

## ROLES OF AMINO ACIDS IN MENTAL HEALTH

Psychoactive substances can be broadly categorised into several groups based on their primary effects on the central nervous system. Stimulants like cocaine and amphetamines elevate mood and increase energy, but they can also lead to anxiety and sleep disorders. Depressants such as alcohol and benzodiazepines have a sedative effect but can cause respiratory issues and dependency. Hallucinogens like LSD and psilocybin alter perception and can induce spiritual experiences, but they may also trigger psychological distress. Opioids like heroin and morphine are potent pain relievers but are highly addictive and can lead to overdoses. Lastly, cannabinoids like marijuana have a range of effects, including relaxation and altered perception, but can also impair memory and coordination. Each of these categories presents unique challenges and considerations in the context of addiction treatment.

Given the multifaceted roles of amino acids in physiological and biochemical functions, they also have significant implications for mental health. For instance, tryptophan serves as a precursor to serotonin [5-hydroxytryptamine (5-HT)], while phenylalanine and tyrosine act as precursors to catecholamine neurotransmitters such as dopamine, norepinephrine, and epinephrine. The synthesis rates of these neurotransmitters in the brain are highly dependent on the availability of their respective dietary precursors (Parker & Brotchie, 2011).

In addition to their primary roles as protein subunits, amino acids like tryptophan and tyrosine offer various other benefits, particularly in mental health. For example, dietary protein enriched in tryptophan has been shown to improve coping ability in stress-vulnerable subjects, likely through alterations in brain serotonin levels (Markus et al., 2000).

Tyrosine, another influential amino acid, serves as a precursor to norepinephrine (NE) and dopamine (DA). Studies in both animals and humans indicate that tyrosine supplementation may mitigate stress-induced NE depletion and improve cognitive performance under stress (Deijen et al., 1999). While the focus here is primarily on tryptophan and tyrosine, other amino acids also exhibit beneficial effects on mood and cognitive function.

## Psychoactive Substances and their Impact on Mental Health

As previously discussed, psychoactive substances can be broadly classified into several categories, each with its

own unique challenges and considerations in the context of addiction treatment.

Psychoactive drugs significantly influence psychological processes such as cognition, perception, and emotion.

### CURRENT TREATMENTS FOR ADDICTION

Substance abuse is characterised by a pattern of compulsive use that leads to significant social, occupational, and interpersonal consequences (APA Dictionary of Psychology). Addiction refers to a state of psychological or physical dependence on alcohol or other drugs, and can also extend to behavioural disorders like internet and gambling addictions (APA Dictionary of Psychology).

Current treatments for addiction and substance abuse typically involve a combination of psychological, behavioural, and pharmacological interventions. Medications such as Naltrexone, Buprenorphine, and Methadone primarily target opioid receptors, while others like Vivitrol affect different brain regions.

Current treatments often come with their own set of limitations, such as potential side effects, high costs, and the risk of developing resistance to or dependency on the medication itself. Moreover, these treatments may not be universally effective, requiring a more personalised approach for different individuals.

Psychoactive substances like cocaine have profound effects on dopaminergic systems, leading to neurotoxicity and other adverse effects (Jitcă et al., 2021). These substances can either mimic, stimulate, or block the effects of neurotransmitters, leading to acute toxicity or chronic issues (Dias da Silva et al., 2021).

There is a strong correlation between neurotransmitter load and oxidative stress, particularly in the context of psychostimulant drug use. These drugs induce the release of monoamines, which are subsequently metabolised, leading to the production of reactive oxygen species (ROS) (Dias da Silva et al., 2021).

### Amino Acids in Mental Health and Addiction Treatment

Beyond the scope of addiction treatment, amino acids have been studied for their efficacy in treating various other mental health conditions. For example, L-tryptophan has been shown to be effective in managing symptoms of depression due to its role as a precursor to serotonin. Similarly, studies on branched-chain amino acids (BCAAs) have indicated their potential in reducing symptoms of mania in bipolar disorder (Fernstrom, 2005). Tyrosine has been researched for its role in improving cognitive functions under stress (Deijen et al., 1999). These studies suggest a broader therapeutic potential for amino acids, making them a subject of interest for comprehensive mental health research.

Amino acids act as the building blocks for neurotransmitters

and play diverse roles in cellular energy balance and stability, according to research by Sato et al. (2020). For example, studies on branched-chain amino acids (BCAAs) have indicated a notable decrease in symptoms of mania in bipolar disorder patients (Fernstrom, 2005).

It is worth noting that amino acids are not just limited to their role in neurotransmitter synthesis; they also participate in other vital physiological processes. For instance, they are essential for protein synthesis, immune function, and even the regulation of blood sugar levels. Their multifaceted roles make them a subject of ongoing research in the fields of neuroscience and metabolism.

Certain amino acids, such as L-tryptophan, have intricate metabolic routes that are crucial for both brain protection and the regulation of the gut-brain axis (Liaqat et al., 2022). In a similar vein, L-tyrosine, which serves as a starting point for catecholamines, has been found to enhance cognitive abilities, particularly under stressful conditions (Bloemendaal et al., 2018).

While the potential benefits of amino acids in treating addiction are promising, it is crucial to note that not all amino acids have been thoroughly studied. Therefore, their use should be approached with caution due to potential side effects (Dias da Silva et al., 2021).

### LIMITATIONS AND FUTURE DIRECTIONS

Despite the promising evidence supporting the role of amino acids in addiction treatment, there are several limitations that need to be addressed. First, not all amino acids have been thoroughly studied for their effects on addiction, leaving gaps in our understanding of their full therapeutic potential. Second, the long-term effects and potential side effects of amino acid supplementation are not yet fully understood, particularly in the context of complex mental health disorders. Future research should aim to fill these gaps by conducting more comprehensive studies, including randomised controlled trials, to ascertain the safety and efficacy of amino acids as an adjunct therapy in addiction treatment.

Moreover, future studies should focus on specific amino acids that have shown preliminary promise but have not been extensively studied in the context of addiction treatment. For instance, the role of amino acids like L-theanine and L-glutamine could be explored further. Additionally, research could also delve into the synergistic effects of combining multiple amino acids in treatment regimens. Longitudinal studies are also needed to assess the long-term safety and efficacy of amino acid supplementation in diverse populations.

### CONCLUSION

The existing literature suggests that amino acids could serve as adjunct treatments in addiction therapy. They play

a crucial role in neurotransmitter production and have neuroprotective effects. However, further research is needed to fully understand their mechanisms of action and potential side effects. Given the promising preliminary evidence, more research is urgently needed to confirm the efficacy of amino acids in addiction treatment. These amino acids have shown promise not only in alleviating symptoms of addiction but also in improving mental health conditions like depression and stress, which often co-occur with substance abuse. The implications of these findings are twofold: First, they suggest that amino acids could be incorporated into a more holistic, patient-centered approach to addiction treatment. Second, they underscore the need for further research to validate these preliminary findings and to explore the mechanisms by which amino acids exert their effects.

## REFERENCES

1. APA Dictionary of Psychology.
2. Bhagavan, N. V., & Ha, C.-E. (2015). Amino Acids. In N. V. Bhagavan & C.-E. Ha (Eds.), *Essentials of Medical Biochemistry* (Second Edition) (pp. 21-29). Academic Press.
3. Bloemendaal, M., Froböse, M. I., Wegman, J., Zandbelt, B. B., van de Rest, O., Cools, R., & Aarts, E. (2018). Neuro-Cognitive Effects of Acute Tyrosine Administration on Reactive and Proactive Response Inhibition in Healthy Older Adults. *Eneuro*, 5(2), ENEURO.0035-17.2018. <https://doi.org/10.1523/eneuro.0035-17.2018>
4. Deijen, J. B., Wientjes, C. J. E., Vullings, H. F. M., Cloin, P. A., & Langefeld, J. J. (1999). Tyrosine improves cognitive performance and reduces blood pressure in cadets after one week of a combat training course. *Brain Research Bulletin*, 48(2), 203–209. [https://doi.org/10.1016/s0361-9230\(98\)00163-4](https://doi.org/10.1016/s0361-9230(98)00163-4)
5. Dias da Silva, D., Silva, J. P., Carmo, H., & Carvalho, F. (2021). Neurotoxicity of psychoactive substances: A mechanistic overview. *Current Opinion in Toxicology*, 28, 76–83. <https://doi.org/10.1016/j.cotox.2021.10.002>
6. Fernstrom, J. D. (2005). Branched-Chain Amino Acids and Brain Function. *The Journal of Nutrition*, 135(6), 1539S-1546S. <https://doi.org/10.1093/jn/135.6.1539S>
7. Jítčá, G., Ůsz, B. E., Tero-Vescan, A., & Vari, C. E. (2021). Psychoactive Drugs—From Chemical Structure to Oxidative Stress Related to Dopaminergic Neurotransmission. A Review. *Antioxidants*, 10(3), 381. <https://doi.org/10.3390/antiox10030381>
8. Liaqat, H., Parveen, A., & Kim, S. Y. (2022). Neuroprotective Natural Products' Regulatory Effects on Depression via Gut–Brain Axis Targeting Tryptophan. *Nutrients*, 14, 3270. <https://doi.org/10.3390/nu14163270>
9. Maloy, S., & Hughes, K. (2013). Amino Acids. In S. Maloy & K. Hughes (Eds.), *Brenner's Encyclopedia of Genetics* (Second Edition) (pp. 108-110). Academic Press.
10. Markus, C. R., Olivier, B., Panhuysen, G. E., Van der Gugten, J., Alles, M. S., Tuiten, A., Westenberg, H. G., Fekkes, D., Koppeschaar, H. F., & de Haan, E. E. (2000). The bovine protein  $\alpha$ -lactalbumin increases the plasma ratio of tryptophan to the other large neutral amino acids, and in vulnerable subjects raises brain serotonin activity, reduces cortisol concentration, and improves mood under stress. *The American Journal of Clinical Nutrition*, 71(6), 1536–1544. <https://doi.org/10.1093/ajcn/71.6.1536>
11. Parker, G., & Brotchie, H. (2011). Mood effects of the amino acids tryptophan and tyrosine. *Acta Psychiatrica Scandinavica*, 124(6), 417–426.

**Citation:** Kian Zehtabian, Dennis Relajo-Howell, et al., “Exploring the Therapeutic Potential of Amino Acids in the Treatment of Psychoactive Substance Addiction: A Comprehensive Review”, *Universal Library of Medical and Health Sciences*, 2023; 1(1): 01-03.

**Copyright:** © 2023 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.