



Comparison between Smoking and Non-Smoking in Term of Thrombocytes Count and Coagulation: Is there any Differences?

Aishe Abdulhafeid Miftah ALgmati¹, Abdalmawla Alhussin Ali Ali², Ashraf Ghaith Mohammed³, Ali Salahuddin Khalifa³, Fatima Ahmed Mohammed³, Fatima Mahjoub ALjuhaimi³, Fatima Shibob Abdullah³, Mohammed Ramadan Farag³

¹Lecturer at Laboratories Department, Faculty of Health Science, Sirte University.

²Professor (Assistant) at Orthodontic Department, Faculty of Dentistry, Sirte University.

³Graduated from Laboratories Department, Faculty of Health Science, Sirte University.

Abstract

Introduction: Smoking is a major risk factor for cardiovascular disease, with a significant impact on platelet activation and thrombosis. This study aims to examine the effects of smoking on platelet activation and INR as a marker of thrombosis. This study aimed to investigate the influence of tobacco smoking on platelets count and coagulation tests (PT, PTT, INR) in adult males in sirte city.

Materials and Methods: A case control study was carried out during 1/1/2023 to 15/2/2023. 200 males (100 smokers and 100 non-smokers) were participated in this study. with ages between 15-85 years enrolled in this study. Data were presented in median (interquartile range). Data was collected by using structural interviewing questionnaire. blood samples were collected in tri sodium citrate anti-coagulant for PT and INR, manually analyzed was using. Ethylene-di-amine-tetra-acetic acid (EDTA) anticoagulant for the platelets count, analyzed by (Sysmex, German) automated hematology analyzer.

Main Results: The results showed that the Median platelets count was statistically no significantly between smokers (237) and non-smokers (256), $U = 5.696.000$, $Z = 1.701$, $P = 0.089$. Whereas, median PT was statistically significant between smokers (15) and non-smokers (14), $U = 4.112.000$, $Z = -2.194$, $P = 0.028$. Also, median INR was statistically significant between smokers (1.4) and non-smokers (1.3), $U = 4.063.500$, $Z = -2.316$, $P = 0.021$.

Conclusion: The present study shows that smokers likely to have lesser platelet counts, longer PT, and higher INR values in comparable to non-smokers. Thus, smoking might be related with bleeding complaints but additional examinations are required.

Keywords: Smoking, Cardiovascular Disease, Thrombocytes Count, Coagulation Test.

INTRODUCTION

Platelets are specialized blood cells formed and released into the bloodstream by precursor cells called megakaryocytes that lie within the bone marrow. They play a critical role in blood clot formation, impacting thrombosis and haemostasis. Because of their capability to become adhesive and release a range of mediators affecting coagulation at the site of injury, they are also involved in immunity, inflammation, and wound healing. Moreover, platelets are directly connected to the coagulation cascade and the humoral immune system through a diverse combination of surface receptors, adhesion proteins, integrins, and glycoproteins that are linked to multiple signalling pathways (Linden, 2013).

The coagulation cascade is a sequence of processes that occur during haemostasis following damage by activating a cascade of proteins known as clotting factors. It has three pathways: intrinsic, extrinsic, and common. The coagulation

pathway's main function is to maintain haemostasis, which is the prevention of bleeding or haemorrhage (Chaudhry *et al*, 2022). Each coagulation pathway comprises coagulation factors that have specific roles in haemostasis. The intrinsic pathway comprises factors I, II, IX, X, XI, and XII. The extrinsic pathway consists of factors I, II, VII, and X. The common pathway consists of factors I, II, V, VIII, X. Coagulation factors are proteins that form blood clots to stop bleeding in the presence of an injury or wound (Chaudhry *et al*, 2022).

Moreover, platelets are critical mediators that initiate the mechanical pathway of the coagulation cascade when blood vessels are damaged. They promote primary haemostasis through three basic mechanisms: activation, adhesion, and aggregation. As the vascular endothelium's integrity is compromised, several macromolecular constituents of the vascular sub-endothelium become exposed and easily accessible to platelets (Periyah *et al*, 2017). Platelets play a significant role in the coagulation cascade by generating

Comparison between Smoking and Non-Smoking in Term of Thrombocytes Count and Coagulation: Is there any Differences?

the initial haemostatic plug, which gives a surface for the synthesis of active coagulation factors, which in turn leads to the development of fibrin stabilized platelet aggregates and subsequent clot retraction (Palta *et al*, 2014).

A variety of factors can influence haemostasis. According to research, cigarette smoking is one of these factors. It can modify the main haemostatic systems and coagulation factors, affecting their functions. Further, smoking has been shown to reduce the expression of tissue factor pathway inhibitors and raise plasma fibrinogen levels in smokers compared to non-smokers (Elkhalifa, 2018). Moreover, the endothelium lining is damaged in smokers, causing platelets to stick to the subendothelial collagen, as a result, the number of platelets in circulation tends to fluctuate (Pujani *et.al*, 2021). According to the results of a study conducted in Nigeria, platelet count was significantly lower in smokers when compared to non-smokers. Additionally, platelet count was lower in older patients compared to younger individuals, and smokers who had smoked for more than ten years had a significant decline in platelet count when compared to those who had smoked for less time. (Okeke *et.al*, 2017).

Smoking may also influence coagulation screening tests, such as blood clotting time. In a study conducted in Sudan aimed to determine the effects of cigarette smoking on coagulation screening tests and platelet counts, the findings revealed that smokers had considerably lower mean of PT and INR compared to those who do not smoke (Elkhalifa, 2018). Moreover, cigarette smoke exposure raises the risk of acute myocardial infarction and sudden cardiac death, with thrombosis being responsible for the majority of these acute occurrences (Barua *et.al*, 2010; Ali AA *et al*, 2023). In addition to affecting haemostasis, numerous studies have connected tobacco use to a variety of fatal diseases such as coronary artery disease, chronic obstructive pulmonary disease, malignancies in every human organ system, and diabetes (Adams & Morris, 2022). Therefore, our study aimed to

investigate the effect of cigarette smoking on platelets count and coagulation tests (PT, PTT, INR) in adult males in Sirte city, Libya.

MATERIALS AND METHODS

Study Population and Methodology

A case control study was carried out during 1/1/2023 to 15/2/2023. 200 males (100 smokers and 100 non-smokers) were participated in this study. with ages between 15-85 years recruited to take place in this study. Data were presented in median (interquartile range). Data was collected by using structural interviewing questionnaire (Appendix.1). Venous blood specimens were collected in tri sodium citrate anti-coagulant for PT and INR, manually analyzed was using. Ethylene-di-amine-tetra-acetic acid (EDTA) anticoagulant for the platelets count, analyzed by (Sysmex, German) automated hematology analyzer.

Statistical Analysis

Data analysis carried out by Statistical Package for the Social Sciences (SPSS) version 25.0 (IBM Corp., Armonk, NY, USA). A Mann-whitne U-test was run to determine if were differences in Platelets count, PT and INR of smokers and non-smokers. Distributions of the engagement scores for smokers and non-smokers were similar, as assessed by visual inspection. Association between weigh lose and smoking status was assessed by Fisher's exact test.

RESULTS

Comparison of Age between Study Groups

200 males are participate in this study. The comparison of median age between smokers and non-smokers group is illustrated in Table 1. As can be seen from Table, the median age of smokers was (27) years, whereas the median age of non-smokers was (30) years. No statistical significance is observe between the age of two groups, $U = 5.414.500$, $Z = 1.014$, $P = 0.311$.

Table 1. Comparison of median age between study groups

Variable	All	Smokers	Non-smokers	P-value
Age	28(23 – 35)	27(23 – 33)	30 (22 – 38)	0.311

The Average of Platelets Count and Coagulation Profile (PT, INR) between Study Groups

Table 2, shows the differences between two groups in platelet counts, PT and INR results. Median platelets count was statistically no significantly between smokers (237) and non-smokers (256), $U = 5.696.000$, $Z = 1.701$, $P = 0.089$. Median PT was statistically significant between smokers (15) and non-smokers (14), $U = 4.112.000$, $Z = -2.194$, $P = 0.028$. Median INR was statistically significant between smokers (1.4) and non-smokers (1.3), $U = 4.063.500$, $Z = -2.316$, $P = 0.021$.

Table 2. Differences in the average of platelet numbers and coagulation profile between the study groups.

Variable	All	Smokers	Non-smokers	P-value
Platelets count	245 (209 – 297)	237 (199 – 278)	256 (216 – 303)	0.089
PT	14 (13 – 16)	15 (13 – 16)	14 (13 – 15.75)	0.028
INR	1.3 (1.2 – 1.5)	1.4 (1.2 – 1.5)	1.3 (1.2 – 1.5)	0.021

Distribution of Diseases between Study Population

As can be seen from Table 3, 168 males (84.0%) have not any disease. While 32 males (16.0%) have diseases such as hypertension, ulcer, respiratory diseases, diabetes and renal diseases.

Table 3, The percentage of diseases between the precipitants

Disease	Frequency	Percent
Hypertension	5	2.5%
Ulcer	6	3.0%
Respiratory diseases	12	6.0%
Diabetes	5	2.5%
Renal diseases	4	2.0%
No diseases	168	84.0%
Total	200	100.0%

Association between smoking status and weigh lose

200 males participated in the study, of the participants, 100 (50%) were smokers and 100 (50%) were non-smokers. For smokers, 36 (83.7%) were have weigh lose, while for non-smokers, 7(16.3%) were have weigh lose. There was statistically significant association between smoking status and weigh lose as assessed by Fisher’s exact test, $P = 0.000$ as show that in table 4.

Table 4. Association between smoking status and weigh lose

Variable	All n(%)	Smokers n (%)	Non-smokers n(%)	P value
Weigh lose	43 (100%)	36(83.7%)	7(16.3%)	0.000

DISCUSSION

Coagulation is the process in which vascular damage is followed by successive platelet adhesion to sites of vessel injury and the initiation of the coagulation cascade. According to research, smoking causes a state of hypercoagulability, which affects the normal platelet physiology and haemostasis pathways (Basalingappa *et al*, 2015). The results of this study shown that the average of platelets count was lower than in smokers compared to non-smokers, however, this difference was not statistically significant. This result is comparable to findings that reported by Asif, *et al* (2013), there was a non-significant decrease in platelets count in smokers compared to those who do not smoke . The drop in platelet count could be the result of smoking damage to the endothelium lining causing platelets to adhere to the sub-endothelial collagen, which in turn causes fluctuations in the amount of platelets in the bloodstream (Pujani, *et al.*, 2021). Nonetheless, our findings disagree with the study that accomplished by Ghahremanfard, *et al* (2015) that reported the smokers had a significantly higher mean platelet count compared to non-smokers.

In the study that accomplished by (Elkhalifa, 2018), it reported that the mean of PT was considerably shorter in smokers compared to non-smokers. In contrast, the results of our study showed that the average of PT was significantly longer in smokers compared to non-smokers. The present study shown that smokers had a greater INR compared to non-smokers. Different findings were reported by Elkhalifa, (2018), the average INR was significantly lower in smokers

compared to the non-smoker group. Data on the impact of smoking on the INR are contradictory (Ali, *et al.*, 2017). However, the majority of studies indicate that smoking reduces INR levels, although the results may differ between each individual (Stading, *et al.*, 2007).

Filozof, *et al.*, (2004) reported that weight change is predicted by younger ages, lower socioeconomic level, and heavier smoking. Regarding the results of the Fisher’s exact test, the findings of the present study demonstrated that there was association between smoking status and weigh lose.

CONCLUSION

The main aim of this research is to investigate how smoking effects on platelets count and coagulation tests. The results indicate that smokers group had a significantly longer PT and INR value, and a slightly decrease platelet counts, in comparison to non-smokers group. Also, there was statistically significant association between smoking status and weigh lose. Therefore, smoking might be related with bleeding complaints but more examinations are required.

REFERENCES

1. Adams, T. N., & Morris, J. (2022). Smoking. In *StatPearls*. StatPearls Publishing.
2. Ali, A. A., Tarawneh, A. M. A., & Abdulqader, M. A. (2017). Level of awareness of maintaining dental records among dental practitioners in Sirte: Are we ready for forensic odontology. *J Dent Dent Med*, 1(2), 2-5.
3. Ali AA, *et al.* (2023). Oral Hygiene Status of Children

Comparison between Smoking and Non-Smoking in Term of Thrombocytes Count and Coagulation: Is there any Differences?

- with Autism Spectrum Disorders: Cross Section Study. *Mathews J Dentistry*. 7(1):34.
4. Asif, Muhammad & Karim, Sajjad & Umar, Zubaida & Malik, Dr & Ismail, Tariq & Chaudhary, Adeel & Al-Qahtani, Mohammed & Rasool, Mahmood. (2013). Effect of cigarette smoking based on hematological parameters: comparison between male smokers and non-smokers. *Turkish Journal of Biochemistry/Turk Biyokimya Dergisi*. 38.
 5. Barua, R. S., Sy, F., Srikanth, S., Huang, G., Javed, U., Buhari, C., Margosan, D., & Ambrose, J. A. (2010). Effects of cigarette smoke exposure on clot dynamics and fibrin structure: an ex vivo investigation. *Arteriosclerosis, thrombosis, and vascular biology*, 30(1), 75–79.
 6. Basalingappa, DoddamaniR & Sandhya, Metta & Satyanarayana, Uppala & Mohanty, Shruti. (2015). Impact of chronic cigarette smoking on platelet aggregation and coagulation profile in apparently healthy male smokers. *International Journal of Clinical and Experimental Physiology*. 2. 128.
 7. Chaudhry, R., Usama, S. M., & Babiker, H. M. (2022). Physiology, Coagulation Pathways. In *StatPearls*. StatPearls Publishing.
 8. Elkhalfa A. M. (2018). Effects of cigarette smoking on coagulation screening tests and platelet counts in a Sudanese male adults population. *Saudi medical journal*, 39(9), 897–901.
 9. Filozof, C., Fernández Pinilla, M. C., & Fernández-Cruz, A. (2004). Smoking cessation and weight gain. *Obesity reviews: an official journal of the International Association for the Study of Obesity*, 5(2), 95–103.
 10. Ghahremanfard, Farahnaz & Semnani, Vahid & Ghorbani, Rouhollah & Malek, Farhad & Behzadfar, Ali & Zahmatkesh, Mehrdad. (2015). Effects of cigarette smoking on morphological features of platelets in healthy men. *Saudi medical journal*. 36. 847-50.
 11. Linden M. D. (2013). Platelet physiology. *Methods in molecular biology (Clifton, N.J.)*, 992, 13–30.
 12. Okeke, Chizoba & EBUKA, EKEANUMBA. (2017). EFFECTS OF CIGARETTE SMOKING ON SOME COAGULATION PARAMETERS OF SMOKERS IN NNEWI METROPOLIS. *Journal of Environmental and Occupational Science*. 6. 12. 10.5455/jeos.20170113084303.
 13. Palta, S., Saroa, R., & Palta, A. (2014). Overview of the coagulation system. *Indian journal of anaesthesia*, 58(5), 515–523.
 14. Periyah, M. H., Halim, A. S., & Mat Saad, A. Z. (2017). Mechanism Action of Platelets and Crucial Blood Coagulation Pathways in Hemostasis. *International journal of hematology-oncology and stem cell research*, 11(4), 319–327.
 15. Pujani, M., Chauhan, V., Singh, K., Rastogi, S., Agarwal, C., & Gera, K. (2021). The effect and correlation of smoking with platelet indices, neutrophil lymphocyte ratio and platelet lymphocyte ratio. *Hematology, transfusion and cell therapy*, 43(4), 424–429.
 16. Stading, J. A., Faulkner, M. A., Skrabal, M. Z., & Chock, A. W. Y. (2007). Effect of tobacco on INR. *American Journal of Health-System Pharmacy*, 64(8), 805. <https://doi.org/10.2146/ajhp070051>

APPENDIX

Personal data

- Age.....
- Are you smoker ?
- How long Smoking period?
- How many cigarettes ?.....

Do you suffer from one the following diseases?

Diseases	Yes	No
Atherosclerosis		
High blood pressure		
Stomach ulcer		
Kidney diseases		
Respiratory or heart diseases		
Are you take blood thinners ?		
Are you suffer from weight loss or gain		

Citation: Aishe Abdulhafeid Miftah ALgmati, Abdalmawla Alhussin Ali Ali, et al., “Comparison between Smoking and Non-Smoking in Term of Thrombocytes Count and Coagulation: Is there any Differences?”, *Universal Library of Medical and Health Sciences*, 2024; 2(2): 01-04. DOI: <https://doi.org/10.70315/uloap.ulmhs.2024.0202001>.

Copyright: © 2024 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.